

NOTES ON A MINIMAL SET OF GENERATORS FOR THE RADICAL IDEAL DEFINING THE DIAGONAL LOCUS OF $(\mathbb{C}^2)^n$

KYUNGYONG LEE AND LI LI

ABSTRACT. We develop several techniques for the study of the radical ideal I defining the diagonal locus of $(\mathbb{C}^2)^n$. Using these techniques, we give combinatorial construction of generators for I of certain bi-degrees.

1. INTRODUCTION

1.1. Overview. Fix a positive integer n . Consider n -tuples of ordered points $\{(x_i, y_i)\}_{1 \leq i \leq n}$ in the plane $\mathbb{C}^2$. The set of all n -tuples forms an affine space $(\mathbb{C}^2)^n$ with coordinate ring $\mathbb{C}[\mathbf{x}, \mathbf{y}] = \mathbb{C}[x_1, y_1, \dots, x_n, y_n]$. The symmetric group S_n acts on $\mathbb{C}[\mathbf{x}, \mathbf{y}]$ by permuting the coordinates in $\mathbf{x}, \mathbf{y}$ simultaneously, that is,

$$\sigma(x_j) := x_{\sigma(j)}, \quad \sigma(y_j) := y_{\sigma(j)} \quad \text{for } \sigma \in S_n.$$

Definition 1. A polynomial $f \in \mathbb{C}[\mathbf{x}, \mathbf{y}]$ is called *alternating* if

$$\sigma(f) = \text{sgn}(\sigma)f \quad \text{for all } \sigma \in S_n.$$

Define $\mathbb{C}[\mathbf{x}, \mathbf{y}]^\epsilon$ to be the vector space of alternating polynomials in $\mathbb{C}[\mathbf{x}, \mathbf{y}]$.

The vector space $\mathbb{C}[\mathbf{x}, \mathbf{y}]^\epsilon$ has a well-known basis which we describe below. Denote by $\mathbb{N}$ the set of nonnegative integers. Let $\mathfrak{D}$ be the set of subsets $D = \{(\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\}$ of $\mathbb{N} \times \mathbb{N}$. For $D \in \mathfrak{D}$, define

$$\Delta(D) := \det \begin{bmatrix} x_1^{\alpha_1} y_1^{\beta_1} & x_1^{\alpha_2} y_1^{\beta_2} & \dots & x_1^{\alpha_n} y_1^{\beta_n} \\ \vdots & \vdots & \ddots & \vdots \\ x_n^{\alpha_1} y_n^{\beta_1} & x_n^{\alpha_2} y_n^{\beta_2} & \dots & x_n^{\alpha_n} y_n^{\beta_n} \end{bmatrix}$$

(by abuse of notation we also use $\Delta(D)$ to denote the above square matrix). Then $\{\Delta(D)\}_{D \in \mathfrak{D}}$ forms a basis for the $\mathbb{C}$ -vector space $\mathbb{C}[\mathbf{x}, \mathbf{y}]^\epsilon$.

The radical ideal I that defines the diagonal locus of $(\mathbb{C}^2)^n$ is

$$I = \bigcap_{1 \leq i < j \leq n} (x_i - x_j, y_i - y_j).$$

A famous theorem of Haiman asserts the following:

Theorem 2. [4, Corollary 3.8.3] *The ideal generated by the alternating polynomials in $\mathbb{C}[\mathbf{x}, \mathbf{y}]$ agrees with I .*

Research of the first author partially supported by NSF grant DMS 0901367.

Haiman's theorem immediately implies that the ideal I is generated by $\{\Delta(D)\}_{D \in \mathfrak{D}}$. He has also proved the following theorem, which asserts that the number of minimal generators of I is equal to the n -th Catalan number.

Theorem 3. ([5, p393]) $\dim_{\mathbb{C}} I/(\mathbf{x}, \mathbf{y})I = \frac{1}{n+1} \binom{2n}{n}$.

Let $M = I/(\mathbf{x}, \mathbf{y})I$. The space M is doubly graded as $M = \oplus_{d_1, d_2} M_{d_1, d_2}$. The t, q -analog of the Catalan number is defined as

$$C_n(q, t) = \sum_{d_1, d_2} t^{d_1} q^{d_2} \dim M_{d_1, d_2}.$$

By a simple algebraic argument, giving a minimal set of generators of I is equivalent to giving a basis of M , and it suffices to find bases for all graded pieces M_{d_1, d_2} . It is then natural to ask the following question:

Question 4. Given a bi-degree (d_1, d_2) , is there a combinatorially significant construction of the basis for each M_{d_1, d_2} ?

On one hand, a combinatorial study of $C_n(q, t)$ gives us hints to construct an explicit basis for M_{d_1, d_2} ; on the other hand, a good understanding of M_{d_1, d_2} helps to study $C_n(q, t)$. For example, if we can show that some of the combinatorially significant elements in M_{d_1, d_2} are linearly independent, then we can give lower or upper bounds for the coefficients appeared in $C_n(q, t)$. This idea is developed further in our subsequent paper [8].

Of course Question 4 is vague and only gives the guideline of study. Here is one way to make it precise, which is one motivation for us to study Question 4. Define

$$\Lambda := \{\lambda = (\lambda_1, \dots, \lambda_n) \mid \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{n-1} \geq \lambda_n = 0, \quad \lambda_i \leq n - i, \forall 1 \leq i \leq n\},$$

$$a(\lambda) := \sum_{i=1}^n (n - i - \lambda_i), \quad \forall \lambda \in \Lambda,$$

$$b(\lambda) := \#\{(i, j) \mid i < j, \lambda_i - \lambda_j + i - j \in \{0, 1\}\}, \quad \forall \lambda \in \Lambda,$$

A surprising combinatorial interpretation for $C_n(q, t)$ found by Garsia and Haglund ([2], [3]) asserts that

$$C_n(q, t) = \sum_{\lambda \in \Lambda} q^{a(\lambda)} t^{b(\lambda)}.$$

It is then natural to ask if there is actually an explicit construction of the basis of M_{d_1, d_2} hidden behind the above combinatorial interpretation. Indeed, a more specified question is posed by Haiman ([6]):

Is there a rule to associate to each $\lambda \in \Lambda$ an element $D(\lambda) \in \mathfrak{D}$ such that $\deg_{\mathbf{x}} \Delta(D(\lambda)) = a(\lambda)$, $\deg_{\mathbf{y}} \Delta(D(\lambda)) = b(\lambda)$, and the set $\{\Delta(D(\lambda))\}_{\lambda \in \Lambda}$ generates I ?

1.2. Techniques and main result. In the study of above questions, we found the following three linear relations that turn the questions into combinatorial games, and lead to a combinatorial construction of bases of M_{d_1, d_2} for certain bi-degrees (d_1, d_2) . First we introduce some notations.

- For $D = \{P_1, \dots, P_n\} \in \mathfrak{D}$ where $P_i = (\alpha_i, \beta_i)$, define $|P_i| = \alpha_i + \beta_i$.
- For two homogeneous polynomials $f, g \in I$ of the same degree d , we denote

$$f \sim g$$


if f and g are equivalent modulo the ideal $I_{<d}$. In other words, under the quotient map $I \rightarrow M = I/(\mathbf{x}, \mathbf{y})I$, the image of f equals the image of g .

Relation 1 (Transfactor Lemma 22). Given positive integers $1 \leq i \neq j \leq n$ such that $|P_i| = i - 1$, $|P_{i+1}| = i$, $|P_j| = j - 1$, $|P_{j+1}| = j$, $\beta_i > 0$, $\alpha_j > 0$ (we assume $|P_{n+1}| = n$). Let D' be obtained from D by moving P_i to southeast and P_j to northwest, i.e.

$$D' = \{P_1, \dots, P_{i-1}, P_i + (1, -1), P_{i+1}, \dots, P_{j-1}, P_j + (-1, 1), P_{j+1}, \dots, P_n\}.$$

Then $\Delta(D) \sim \Delta(D')$.

Example: $n = 9, i = 2, j = 6$.




Relation 2 (Permuting Lemma 24). Given positive integers h, ℓ and m such that $2 \leq h < h + \ell + m \leq n + 1$, $|P_h| = h - 1$, $|P_{h+\ell}| = h + \ell - 1$, $|P_{h+\ell+m}| = h + \ell + m - 1$ (by convention, the last equality holds if $h + \ell + m = n + 1$) and $\alpha_{h+\ell}, \dots, \alpha_{h+\ell+m-1} \geq \ell$. Let D' be obtained from D by moving the m points $P_{h+\ell}, \dots, P_{h+\ell+m-1}$ to the left by ℓ units and moving the ℓ points $P_h, \dots, P_{h+\ell-1}$ to the right by m units, i.e.

$$D' = \{P_1, P_2, \dots, P_{h-1}, P_{h+\ell} - (\ell, 0), P_{h+\ell+1} - (\ell, 0), \dots, P_{h+\ell+m-1} - (\ell, 0), \\ P_h + (m, 0), P_{h+1} + (m, 0), \dots, P_{h+\ell-1} + (m, 0), P_{h+\ell+m}, \dots, P_n\}.$$

Then $\Delta(D) \sim \Delta(D')$.

Example: $n = 10, h = 3, \ell = 4, m = 3$.




Relation 3 (Lemma 29). Given positive integers j and s . Suppose P_{s_0} is the last point in D satisfying $|P_i| = i - 1$. Define $j = (s_0 - 1 - |P_{s_0}|) + (s_0 - |P_{s_0+1}|) + \dots + (n - 1 - |P_n|)$. Suppose $|P_i| = i - 1$ for $1 \leq i \leq j + 2$, $P_2 = (1, 0)$, $s_0 \leq s \leq n$, and $\alpha_s, \beta_s \geq 1$. Let

$$D^{\nwarrow} = \{P_1, \dots, P_{j+1}, P_{j+2} + (1, -1), P_{j+3}, \dots, P_{s-1}, P_s + (-1, 1), P_{s+1}, \dots, P_n\}, \\ D^{\searrow} = \{P_1, (0, 1), P_3, \dots, P_{s-1}, P_s + (1, -1), P_{s+2}, \dots, P_n\}.$$

Then $2\Delta(D) \sim \Delta(D^{\nwarrow}) + \Delta(D^{\searrow})$.

Example: $n = 9, i = 2, j = 6$.




By playing with the above three relations, we can easily obtain our main theorem. Let us first define minimal staircase forms.

Definition 5. We call $D = \{P_1, \dots, P_n\}$ a *minimal staircase form* if $|P_i| = i - 1$ or $i - 2$ for every $1 \leq i \leq n$. For a minimal staircase form D , let $\{i_1 < i_2 < \dots < i_\ell\}$ be the set of i 's such that $|P_i| = i - 1$, we define the *partition type* of D to be the partition of $(\binom{n}{2} - \sum |P_i|)$ consisting of all the positive integers in the sequence

$$(i_1 - 1, i_2 - i_1 - 1, i_3 - i_2 - 1, \dots, i_\ell - i_{\ell-1} - 1, n - i_\ell).$$

Example: Let $n = 8$ and $D = \{P_1, \dots, P_8\}$ satisfying $(|P_1|, \dots, |P_8|) = (0, 1, 1, 2, 4, 4, 5, 6)$. Then D is a minimal staircase form. The set $\{i \mid |P_i| = i - 1\}$ equals $\{1, 2, 5\}$. The positive integers in the sequence $(1 - 1, 2 - 1 - 1, 5 - 2 - 1, 8 - 5)$ are $(2, 3)$, so the partition type of D is $(2, 3)$.



Let $p(k)$ denote the number of partitions of an integer k and Π_k denote the set of partitions of k .

Theorem 6 (Main Theorem). *Let k be any positive integer. There are positive constants $c_1 = 8k + 5, c_2 = 2k + 1$ such that the following holds:*

For integers n, d_1, d_2 satisfying $n \geq c_1, d_1 \geq c_2 n, d_2 \geq c_2 n$ and $d_1 + d_2 = \binom{n}{2} - k$, the vector space M_{d_1, d_2} has dimension $p(k)$, and the $p(k)$ elements

$$\{ \text{a minimal staircase form of bi-degree } (d_1, d_2) \text{ and of partition type } \mu \}_{\mu \in \Pi_k}$$

form a basis of M_{d_1, d_2} .

Remark 7. Bergeron and Chen have found explicit bases for M_{d_1, d_2} for certain bi-degrees using a different method [1].

Let $p(\delta, k)$ be the number of partitions of k into at most δ parts. It is elementary that $p(\delta, k)$ is the same as the number of partitions of k into parts no larger than δ (for example, [7, p.83]). By convention $p(\delta, 0) = 1$ for $\delta \geq 0$; $p(0, k) = 0$ for $k > 0$. We pose the following conjecture generalizing the main theorem.

Conjecture 8. Let $k \leq n - 3$ be a non-negative integer. Let d_1, d_2 be two non-negative integers such that $d_1 + d_2 = n(n - 1)/2 - k$. Let $\delta = \min\{d_1, d_2\}$. Then

$$\dim M_{d_1, d_2} = p(\delta, k),$$

and there is a basis that can be constructed combinatorially.

Remark 9. The conjecture is proved in our subsequent paper [8].

The structure of the paper is as follows. In §2 we give the definition of staircase forms and discuss their properties. In §3 we prove the three relations given at the beginning of §1.2, and at the end of this section we give the proof of the main theorem. §4 gives a conjectural minimal set of generators as an answer to Haiman's question, which is equivalent to a conjecture of Mahir Can and Nick Loehr.

Acknowledgements. The authors thank Jim Haglund, Alexander Woo and Alexander Yong for many suggestions and shared insights. We thank the anonymous referees for helpful and constructive suggestions and comments on the paper.

2. ASYMPTOTIC BEHAVIOR OF t, q -CATALAN NUMBERS

In this section, we first introduce the notion of staircase form and block diagonal form, which are matrices whose determinants are equivalent to $\Delta(D)$ modulo $(\mathbf{x}, \mathbf{y})I$. Then we define the partition type of a staircase form. Finally we give Corollary 21, which is half of the main theorem.

Definition 10. Let $D = \{P_1, \dots, P_n\}$ where $P_i = (\alpha_i, \beta_i)$ for $1 \leq i \leq n$. Define $s_i := \alpha_i + \beta_i$. We say D is *in standard order*, if (i) $s_1 \leq s_2 \leq \dots \leq s_n$, and (ii) if $s_i = s_{i+1}$ then $\alpha_i < \alpha_{i+1}$.

Definition-Proposition 11 (Staircase form). Let $D = \{(\alpha_1, \beta_1), (\alpha_2, \beta_2), \dots, (\alpha_n, \beta_n)\} \in \mathfrak{D}$. Define $s_j := \alpha_j + \beta_j$. Define $d := \sum_j s_j = \sum_j (\alpha_j + \beta_j)$ the degree of D , and $(d_1, d_2) := (\sum \alpha_j, \sum \beta_j)$ the bi-degree of D . Define $k = n(n - 1)/2 - d$ the *deficit* of D . Denote by $I_{<d}$ the ideal of $\mathbb{C}[\mathbf{x}, \mathbf{y}]$ generated by homogeneous elements of degree less than d in I . Then there is a matrix S whose (i, j) -th entry is

$$\begin{cases} 0, & \text{if } i \leq s_j; \\ a_{i1}a_{i2} \cdots a_{i, s_j} \text{ where } a_{i\ell} \text{ is either } x_i - x_\ell \text{ or } y_i - y_\ell, & \text{otherwise,} \end{cases}$$

for all $1 \leq i, j \leq n$, such that $\det S \sim \Delta(D)$ modulo $I_{<d}$. Rearranging the columns of S if necessary, we assume that $s_1 \leq s_2 \leq \dots \leq s_n$. We call the matrix S , or its determinant $\det S$, a *staircase form* of D .

Proof. The idea is to construct a matrix that is as close as possible to an Echelon form modulo $I_{<d}$.

For simplicity of notation, denote $x_{ij} = x_i - x_j$, $y_{ij} = y_i - y_j$. If $\alpha_1 > 0$, then the first column of the matrix $\Delta(D)$

$$[x_1^{\alpha_1} y_1^{\beta_1}, \dots, x_n^{\alpha_1} y_n^{\beta_1}]^T$$

equals to

$$x_1[x_1^{\alpha_1-1} y_1^{\beta_1}, \dots, x_n^{\alpha_1-1} y_n^{\beta_1}]^T + [0, x_2^{\alpha_1-1} x_{21} y_2^{\beta_1}, \dots, x_n^{\alpha_1-1} x_{n1} y_n^{\beta_1}]^T.$$

Therefore

$$\Delta(D) = x_1 \det \begin{bmatrix} x_1^{\alpha_1-1} y_1^{\beta_1} & \cdots & x_1^{\alpha_n} y_1^{\beta_n} \\ \vdots & \ddots & \vdots \\ x_n^{\alpha_1-1} y_n^{\beta_1} & \cdots & x_n^{\alpha_n} y_n^{\beta_n} \end{bmatrix} + \det \begin{bmatrix} 0 & x_1^{\alpha_2} y_1^{\beta_2} & \cdots & x_1^{\alpha_n} y_1^{\beta_n} \\ x_2^{\alpha_1-1} x_{21} y_2^{\beta_1} & x_2^{\alpha_2} y_2^{\beta_2} & \cdots & x_2^{\alpha_n} y_2^{\beta_n} \\ \vdots & \vdots & \ddots & \vdots \\ x_n^{\alpha_1-1} x_{n1} y_n^{\beta_1} & x_n^{\alpha_2} y_n^{\beta_2} & \cdots & x_n^{\alpha_n} y_n^{\beta_n} \end{bmatrix}$$

The first summand is a polynomial in $I_{<d}$, so $\Delta(D)$ is equivalent to the second summand modulo $I_{<d}$. If $\alpha_1 - 1 > 0$, we write the first column of the second matrix

$$[0, x_2^{\alpha_1-1} x_{21} y_2^{\beta_1}, \dots, x_n^{\alpha_1-1} x_{n1} y_n^{\beta_1}]^T$$

as a sum of two vectors

$$x_2[0, x_2^{\alpha_1-2} x_{21} y_2^{\beta_1}, \dots, x_n^{\alpha_1-2} x_{n1} y_n^{\beta_1}]^T + [0, 0, x_3^{\alpha_1-2} x_{32} x_{31} y_3^{\beta_1}, \dots, x_n^{\alpha_1-2} x_{n2} x_{n1} y_n^{\beta_1}]^T.$$

Then by a similar argument as above, $\Delta(D)$ is equivalent to

$$\det \begin{bmatrix} 0 & x_1^{\alpha_2} y_1^{\beta_2} & \cdots & x_1^{\alpha_n} y_1^{\beta_n} \\ 0 & x_2^{\alpha_2} y_2^{\beta_2} & \cdots & x_2^{\alpha_n} y_2^{\beta_n} \\ x_3^{\alpha_1-2} x_{32} x_{31} y_3^{\beta_1} & x_3^{\alpha_2} y_3^{\beta_2} & \cdots & x_3^{\alpha_n} y_3^{\beta_n} \\ \vdots & \vdots & \ddots & \vdots \\ x_n^{\alpha_1-2} x_{n2} x_{n1} y_n^{\beta_1} & x_n^{\alpha_2} y_n^{\beta_2} & \cdots & x_n^{\alpha_n} y_n^{\beta_n} \end{bmatrix}$$

modulo $I_{<d}$. If $\beta_1 > 0$, we can apply the similar operation. Repeating this operation, we will eventually replace the first column by the following column vector

$$\begin{bmatrix} 0 \\ \vdots \\ 0 \\ x_{s_1+1,1} x_{s_1+1,2} \cdots x_{s_1+1,\alpha_1} y_{s_1+1,\alpha_1+1} y_{s_1+1,\alpha_1+2} \cdots y_{s_1+1,s_1} \\ x_{s_1+2,1} x_{s_1+2,2} \cdots x_{s_1+2,\alpha_1} y_{s_1+2,\alpha_1+1} y_{s_1+2,\alpha_1+2} \cdots y_{s_1+2,s_1} \\ \vdots \\ x_{n1} x_{n2} \cdots x_{n,\alpha_1} y_{n,\alpha_1+1} y_{n,\alpha_1+2} \cdots y_{n,s_1} \end{bmatrix}$$

where the first $\min\{s_1, n\}$ entries are 0. Note that we may use a different order of operations with respect to x_i or y_i , and the nonzero entries in the first column result might be different.

Applying this procedure for every column, we get a matrix with $\min\{s_j, n\}$ zeros at the j -th column for $1 \leq j \leq n$. Rearrange the columns such that the numbers of zeros in the columns are weakly increasing from left to right. The resulting matrix is a staircase form S of D . $\square$

Corollary 12. *Let D and S be defined as in Definition-Proposition 11. If $s_j > j - 1$ for some $1 \leq j \leq n$, then $\Delta(D) \in I_{<d}$.*

Proof. It is easy to see that $\det S = 0$. $\square$

Definition 13. Let D and S be defined as in Definition-Proposition 11. Consider the set $\{j : s_j = j - 1\} = \{r_1 < r_2 < \cdots < r_\ell\}$ and define $r_{\ell+1} = n + 1$. For $1 \leq t \leq \ell$, define the t -th block B_t of S to be the square submatrix of S of size $(r_{t+1} - r_t)$ whose upper-left

corner is the (r_t, r_t) -entry. Define the *block diagonal form* $B(S)$ of S to be the block diagonal matrix $\text{diag}(B_1, \dots, B_\ell)$.

It is easy to see that $\det B(S) = \det S$.

Example 14. Let $D = \{(0, 0), (1, 0), (0, 2), (1, 1), (3, 1)\}$. It is in standard order. Then

$$\Delta(D) = \begin{bmatrix} 1 & x_1 & y_1^2 & x_1 y_1 & x_1^3 y_1 \\ 1 & x_2 & y_2^2 & x_2 y_2 & x_2^3 y_2 \\ 1 & x_3 & y_3^2 & x_3 y_3 & x_3^3 y_3 \\ 1 & x_4 & y_4^2 & x_4 y_4 & x_4^3 y_4 \\ 1 & x_5 & y_5^2 & x_5 y_5 & x_5^3 y_5 \end{bmatrix},$$

and one possible staircase form of D is

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & x_{21} & 0 & 0 & 0 \\ 1 & x_{31} & y_{31} y_{32} & x_{31} y_{32} & 0 \\ 1 & x_{41} & y_{41} y_{42} & x_{41} y_{42} & 0 \\ 1 & x_{51} & y_{51} y_{52} & x_{51} y_{52} & x_{51} y_{52} x_{53} x_{54} \end{bmatrix}.$$

The corresponding block diagonal form is

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & x_{21} & 0 & 0 & 0 \\ 0 & 0 & y_{31} y_{32} & x_{31} y_{32} & 0 \\ 0 & 0 & y_{41} y_{42} & x_{41} y_{42} & 0 \\ 0 & 0 & 0 & 0 & x_{51} y_{52} x_{53} x_{54} \end{bmatrix}.$$

□

Now we give an alternative definition of minimal staircase form and its partition type. It is equivalent to Definition 5.

Definition 15. Suppose that $\mu = \sum m_i j_i \in \Pi_k$ is a partition of k , where j_i are distinct positive integers. Given a nonzero staircase form S , if for each i the block diagonal form $B(S)$ contains exactly m_i blocks with each having j_i nonzero entries above the diagonal, then we say S is of *partition type* μ . Furthermore, if

$$(2.1) \quad (\text{the entry in the } i\text{-th row and } j\text{-th column in } S) = 0 \text{ for every } i, j \text{ with } j > i + 1,$$

then S is called a *minimal* staircase form of partition type μ . We call a block is minimal if the block satisfies condition (2.1). □

Example 16. Let $n = 11$, $k = 7$, $s_1 = 0$, $s_2 = 1$, $s_3 = 2$, $s_4 = 2$, $s_5 = 4$, $s_6 = 4$, $s_7 = 4$, $s_8 = 7$, $s_9 = 7$, $s_{10} = 8$ and $s_{11} = 9$. Then S is a staircase form of partition type $3 + 3 + 1$, but is not minimal because there is a nonzero entry in the fifth row and seventh column. The 4-th block is not minimal.

$$S = \begin{bmatrix} * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ * & * & * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ * & * & * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ * & * & * & * & * & * & 0 & 0 & 0 & 0 & 0 \\ * & * & * & * & * & * & * & 0 & 0 & 0 & 0 \\ * & * & * & * & * & * & * & * & 0 & 0 & 0 \\ * & * & * & * & * & * & * & * & * & 0 & 0 \\ * & * & * & * & * & * & * & * & * & * & 0 \\ * & * & * & * & * & * & * & * & * & * & * \\ * & * & * & * & * & * & * & * & * & * & * \end{bmatrix}, \quad B(S) = \begin{bmatrix} * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & * & * & * & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & * & * & * & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & * & * & * & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & * & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & * & * \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & * & * \end{bmatrix}. \quad \square$$

Definition 17. Define a natural partial order on the set of partitions Π_k as follows: for two partitions $\mu = (\mu_1 + \cdots + \mu_s)$ and $\nu = (\nu_1 + \cdots + \nu_t)$ in Π_k , define $\mu <_P \nu$ if $\mu \neq \nu$ and μ is a subpartition of ν , i.e., if it is possible to rearrange the order of μ as $(\mu'_1 + \cdots + \mu'_s)$ such that there exist $0 = i_0 < i_1 < \cdots < i_{t-1} < i_t = s$ satisfying

$$\mu'_{i_{r-1}+1} + \mu'_{i_{r-1}+2} + \cdots + \mu'_{i_r} = \nu_r, \quad \text{for } r = 1, 2, \dots, t.$$

Define $\mu \leq_P \nu$ if $\mu = \nu$ or $\mu <_P \nu$. $\square$

Example 18. In the set Π_{10} of partitions of 10, we have $(4 + 2 + 2 + 1 + 1) <_P (5 + 3 + 2)$ because we can rearrange $(4 + 2 + 2 + 1 + 1)$ to $(4 + 1 + 2 + 1 + 2)$, and $4 + 1 = 5$, $2 + 1 = 3$, $2 = 2$. $\square$

The following two propositions are essential ingredients to prove the main theorem. We will only state the propositions here but leave the proofs to §3.

Proposition 19. Suppose $n \geq 8k+5$, $d_1, d_2 \geq (2k+1)n$ and fix a partition $\mu = \sum m_i j_i \in \Pi_k$. Then any nonzero staircase form f of type μ of bidegree (d_1, d_2) is in the ideal

$$I_{<d} + (\text{minimal staircase forms of bidegree } (d_1, d_2) \text{ and of partition types } \leq_P \mu),$$

that is to say, f can be generated by elements in $I_{<d}$ and minimal staircase forms of the same or lower partition types of bidegree (d_1, d_2) .

Proposition 20. Suppose $n \geq 8k+5$, $d_1, d_2 \geq (2k+1)n$ and fix a partition $\mu = \sum m_i j_i \in \Pi_k$. Then any minimal staircase form of partition type μ generates all the minimal staircase forms of the same partition type μ , modulo the ideal

$$I_{<d} + (\text{minimal staircase forms of partition types } <_P \mu).$$

Corollary 21. Suppose $n \geq 8k + 5$, $d_1, d_2 \geq (2k + 1)n$. Then M_{d_1, d_2} can be generated by $p(k)$ elements $\{\det(S_\mu)\}_{\mu \in \Pi_k}$, where for each partition $\mu \in \Pi_k$, S_μ is an arbitrary minimal staircase form of bidegree (d_1, d_2) and of partition type μ . In particular, $\dim M_{d_1, d_2} \leq p(k)$.

Proof. It is an immediate consequence of Proposition 19 and Proposition 20. $\square$

3. PROOF OF MAIN THEOREM

This section is the most technical part of the paper. First we give Transfactor Lemma (Lemma 22) and Minors Permuting Lemma (Lemma 24), which are simple but powerful tools to modify $D \in \mathfrak{D}$ of degree d to another $D' \in \mathfrak{D}$ such that $\Delta(D) \sim \Delta(D')$ modulo $I_{<d}$. Then we prove Lemma 29 which gives a relation among the determinants of D and

certain modifications of D . After this lemma is established, we shall prove Proposition 20, Proposition 19 and then the main theorem (Theorem 6).

Lemma 22 (Transfactor Lemma). *Let $D = \{P_1, \dots, P_n\} \in \mathfrak{D}$ where $P_i = (\alpha_i, \beta_i) \in \mathbb{N} \times \mathbb{N}$ and define $\deg P_i := s_i (= \alpha_i + \beta_i)$ for $1 \leq i \leq n$ and $d = \sum s_i$. Define $s_{n+1} = n$. Suppose $1 \leq i \neq j \leq n$ are two integers satisfying $s_i = i - 1$, $s_{i+1} = i$, $s_j = j - 1$, $s_{j+1} = j$, $\beta_i > 0$, $\alpha_j > 0$. Define*

$$D' = \{P_1, \dots, P_{i-1}, P_i + (1, -1), P_{i+1}, \dots, P_{j-1}, P_j + (-1, 1), P_{j+1}, \dots, P_n\}.$$

Then $\Delta(D) \sim \Delta(D')$ modulo $I_{<d}$.

Proof. By performing appropriate operations as in Definition-Proposition 11, we can obtain a staircase form S of D (resp. staircase form S' of D'), such that the (i, i) -entry and (j, j) -entry of S (resp. S') are $y_{i1} \prod_{t=2}^{i-1} a_{it}$ and $x_{j1} \prod_{t=2}^{j-1} a_{jt}$ (resp. $x_{i1} \prod_{t=2}^{i-1} a_{it}$ and $y_{j1} \prod_{t=2}^{j-1} a_{jt}$). The block diagonal forms of S and S' only differ at two blocks of size 1 located at the (i, i) -entry and (j, j) -entry. Let f_0 be the product of determinants of all blocks of $B(S)$ except the (i, i) -entry and (j, j) -entry. Then $\Delta(D) - \Delta(D')$ is equivalent to the following modulo $I_{<d}$,

$$\begin{aligned} \det(S) - \det(S') &= \left(y_{i1} \prod_{t=2}^{i-1} a_{it} \right) \left(x_{j1} \prod_{t=2}^{j-1} a_{jt} \right) f_0 - \left(x_{i1} \prod_{t=2}^{i-1} a_{it} \right) \left(y_{j1} \prod_{t=2}^{j-1} a_{jt} \right) f_0 \\ &= -\det \begin{bmatrix} 1 & x_1 & y_1 \\ 1 & x_i & y_i \\ 1 & x_j & y_j \end{bmatrix} \left(\prod_{t=2}^{i-1} a_{it} \right) \left(\prod_{t=2}^{j-1} a_{jt} \right) f_0. \end{aligned}$$

Without loss of generality, assume $i < j$. Since

$$I = \bigcap_{1 \leq i < j \leq n} (x_i - x_j, y_i - y_j),$$

it is easy to see that $(\det(S) - \det(S'))/a_{ji}$ is a polynomial in $I_{<d}$ and then the lemma follows. $\square$

The Transfactor Lemma immediately leads to the proof of the following lemma, which is the base case $k = 0$ of the inductive proof of Proposition 20.

Lemma 23. *Let d_1, d_2 be two non-negative integers such that $d_1 + d_2 = n(n-1)/2$. Then M_{d_1, d_2} is generated by any single nonzero staircase form modulo $I_{<n(n-1)/2}$.*

Proof. Let S be a staircase form with $\det S \neq 0$ and bidegree (d_1, d_2) . Because $d_1 + d_2 = n(n-1)/2$, there are $n(n-1)/2$ zeros in the staircase form S . Since $\det S \neq 0$, S and its block diagonal form $B(S)$ must be of the following form

$$S = \begin{bmatrix} * & 0 & \cdots & 0 & 0 \\ * & * & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ * & * & \cdots & * & 0 \\ * & * & \cdots & * & * \end{bmatrix}, \quad B(S) = \begin{bmatrix} * & 0 & \cdots & 0 & 0 \\ 0 & * & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & * & 0 \\ 0 & 0 & \cdots & 0 & * \end{bmatrix}.$$

By repeatedly applying Transfactor Lemma we can easily deduce the following assertion: if S' is a staircase form of another D' of the same bidegree (d_1, d_2) as D , then $\det B(S') \sim \det B(S)$ modulo $I_{<n(n-1)/2}$. The lemma follows from this assertion. $\square$

Lemma 24 (Minors Permuting Lemma). *Let $D = \{P_1, \dots, P_n\} \in \mathfrak{D}$ where $P_i = (\alpha_i, \beta_i)$. Suppose h, ℓ and m are positive integers satisfying $2 \leq h < h + \ell + m \leq n + 1$, $s_h = h - 1$, $s_{h+\ell} = h + \ell - 1$, $s_{h+\ell+m} = h + \ell + m - 1$ (this condition holds if $h + \ell + m = n + 1$ since we assume $s_{n+1} = n$) and suppose that $\alpha_{h+\ell}, \dots, \alpha_{h+\ell+m-1} \geq \ell$. Define*

$$D' = \{P_1, P_2, \dots, P_{h-1}, P_{h+\ell} - (\ell, 0), P_{h+\ell+1} - (\ell, 0), \dots, P_{h+\ell+m-1} - (\ell, 0), \\ P_h + (m, 0), P_{h+1} + (m, 0), \dots, P_{h+\ell-1} + (m, 0), P_{h+\ell+m}, \dots, P_n\}.$$

Then $\Delta(D) \sim \Delta(D')$ modulo $I_{<d}$.

Proof. By performing appropriate operations as in Definition-Proposition 11 and using the assumption that $\alpha_v \geq \ell$ for $h + \ell \leq v \leq h + \ell + m - 1$, we can obtain a staircase form S of D , where the (u, v) -entry for $h + \ell \leq u, v \leq h + \ell + m - 1$ contains the factor $\prod_{j=h}^{h+\ell-1} x_{uj} = \prod_{j=h}^{h+\ell-1} (x_u - x_j)$. Let $B(S) = \text{diag}(B_1, B_2, \dots, B_s)$ be the block diagonal form of S , and let B_r (resp. B_{r+1}) be the block of size ℓ (resp. m) whose upper left corner is the (h, h) -entry (resp. $(h + \ell, h + \ell)$ -entry). Then by our choice of S , all entries in the i -th row ($1 \leq i \leq m$) of B_{r+1} contain $\prod_{j=h}^{h+\ell-1} x_{i+h+\ell-1,j}$ as a factor. Dividing the i -th row of B_{r+1} by $\prod_{j=h}^{h+\ell-1} x_{i+h+\ell-1,j}$ for $1 \leq i \leq m$ and multiplying the i' -th row of B_r by $\prod_{j=h+\ell}^{h+\ell+m-1} x_{i'+h-1,j}$ for $1 \leq i' \leq \ell$, we obtain a new block diagonal matrix $B' = \text{diag}(B_1, \dots, B_{r-1}, B'_r, B'_{r+1}, B_{r+2}, \dots, B_s)$. Since

$$\prod_{j=h}^{h+\ell-1} x_{i+h+\ell-1,j} = (-1)^{\ell m} \prod_{j=h}^{h+\ell-1} x_{i+h+\ell-1,j},$$

$$(-1)^{\ell m} \det B' = \det B = \det S.$$

Now interchange the two blocks B'_r and B'_{r+1} in B' and then change the indices $1, \dots, n$ to

$$1, \dots, (\ell - 1), (\ell + h), \dots, (\ell + h + m - 1), \ell, \dots, (\ell + h - 1), (\ell + h + m), \dots, n.$$

The resulting matrix is the block diagonal matrix of a staircase form of D' . Notice that when we change the indices, the determinant of the resulting matrix is equal to $\det B'$ multiplied by $(-1)^{\ell m}$. Therefore $\Delta(D)$ and $\Delta(D')$ are equivalent modulo $I_{<d}$. $\square$

Example 25. In Example 16, assume $\alpha_8, \dots, \alpha_{11} \geq 3$. Lemma 24 asserts that by permuting the two blocks (as framed in the following figure) in the block diagonal form, the determinant is not changed modulo $I_{<d}$. That is to say, we may permute adjacent blocks provided that the α_i 's in the second block is not less than the size of the first block.

$$\begin{bmatrix}
* & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & * & * & * & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & * & * & * & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & * & * & * & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & * & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & * & * & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & * & * & *
\end{bmatrix}
\longrightarrow
\begin{bmatrix}
* & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & * & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & * & * & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & * & * & * & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & * & * & * & * & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & * & * \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & * & * \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & * & * & *
\end{bmatrix}$$

We frequently use the following elementary lemma.

Lemma 26. *For any non-negative integers c and e ,*

$$\begin{aligned}
& (x_1^c y_1^e + \cdots + x_n^c y_n^e) \cdot \det \begin{bmatrix} x_1^{\alpha_1} y_1^{\beta_1} & x_1^{\alpha_2} y_1^{\beta_2} & \cdots & x_1^{\alpha_n} y_1^{\beta_n} \\ x_2^{\alpha_1} y_2^{\beta_1} & x_2^{\alpha_2} y_2^{\beta_2} & \cdots & x_2^{\alpha_n} y_2^{\beta_n} \\ \vdots & \vdots & \ddots & \vdots \\ x_n^{\alpha_1} y_n^{\beta_1} & x_n^{\alpha_2} y_n^{\beta_2} & \cdots & x_n^{\alpha_n} y_n^{\beta_n} \end{bmatrix} \\
&= \det \begin{bmatrix} x_1^{\alpha_1+c} y_1^{\beta_1+e} & x_1^{\alpha_2} y_1^{\beta_2} & \cdots & x_1^{\alpha_n} y_1^{\beta_n} \\ x_2^{\alpha_1+c} y_2^{\beta_1+e} & x_2^{\alpha_2} y_2^{\beta_2} & \cdots & x_2^{\alpha_n} y_2^{\beta_n} \\ \vdots & \vdots & \ddots & \vdots \\ x_n^{\alpha_1+c} y_n^{\beta_1+e} & x_n^{\alpha_2} y_n^{\beta_2} & \cdots & x_n^{\alpha_n} y_n^{\beta_n} \end{bmatrix} + \cdots + \det \begin{bmatrix} x_1^{\alpha_1} y_1^{\beta_1} & x_1^{\alpha_2} y_1^{\beta_2} & \cdots & x_1^{\alpha_n+c} y_1^{\beta_n+e} \\ x_2^{\alpha_1} y_2^{\beta_1} & x_2^{\alpha_2} y_2^{\beta_2} & \cdots & x_2^{\alpha_n+c} y_2^{\beta_n+e} \\ \vdots & \vdots & \ddots & \vdots \\ x_n^{\alpha_1} y_n^{\beta_1} & x_n^{\alpha_2} y_n^{\beta_2} & \cdots & x_n^{\alpha_n+c} y_n^{\beta_n+e} \end{bmatrix}.
\end{aligned}$$

Proof. The Lemma is the special case of the following Lemma where

$$p = x_1^c y_1^e, \quad q = x_1^{\alpha_1} y_1^{\beta_1} x_2^{\alpha_2} y_2^{\beta_2} \cdots x_n^{\alpha_n} y_n^{\beta_n}.$$

□

Lemma 27. *For $p, q \in \mathbb{C}[\mathbf{x}, \mathbf{y}]$, we have*

$$A(\text{Sym}(p)q) = \text{Sym}(p)A(q)$$

where $\text{Sym}(p)$ denotes the symmetric sum $\sum_{\sigma \in S_n} \sigma(p)$ and $A(p)$ denotes the alternating sum $\sum_{\sigma \in S_n} \text{sign}(\sigma) \sigma(p)$.

Proof. Since the polynomial $\text{Sym}(p)$ is invariant under S_n action,

$$A(\text{Sym}(p)q) = \sum_{\sigma} \text{sign}(\sigma) \sigma(\text{Sym}(p)q) = \sum_{\sigma} \text{sign}(\sigma) \text{Sym}(p) \sigma(q) = \text{Sym}(p)A(q).$$

□

Remark 28. Lemma 26 implies that the determinants on the right hand side of the equality are linearly dependent modulo $I_{<d}$ where $d = \sum_{i=1}^n (\alpha_i + \beta_i) + c + e$. It is easier to express the dependency in terms of squares in $\mathbb{N} \times \mathbb{N}$: for $D = \{(\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\} \in \mathfrak{D}$, let $D_i \in \mathfrak{D}$ be obtained from D by replacing (α_i, β_i) by $(\alpha_i + c, \beta_i + e)$. Then Lemma 26 asserts that

$$\Delta(D_1) + \Delta(D_2) + \cdots + \Delta(D_n) \sim 0 \quad \text{modulo } I_{<d}.$$

Up to modulo $I_{<d}$, we can replace $\Delta(D_i)$ by a linear combination of $\Delta(D_j)$ for $j \neq i$. To say it more vividly, D_j is obtained from D_i by sending $(\alpha_i + c, \beta_i + e)$ to (α_i, β_i) , and then sending (α_j, β_j) to $(\alpha_j + c, \beta_j + e)$. $\square$

Lemma 29. *Let $D = \{P_1, \dots, P_n\} \in \mathfrak{D}$ where $P_i = (\alpha_i, \beta_i)$ are not necessarily distinct and $\{s_i := \alpha_i + \beta_i\}_{1 \leq i \leq n}$ are weakly increasing. Let S be a staircase form of D and $B(S)$ its block diagonal form. Suppose the last block of $B(S)$ is of size t_0 and in this block there are j_r nonzero entries above the diagonal. Suppose the first $(j_r + 2)$ blocks of $B(S)$ are of size 1, i.e., $s_i = i - 1$ for $1 \leq i \leq j_r + 3$. Suppose $P_2 = (1, 0)$. Let t be an integer that $1 \leq t \leq t_0$. Suppose $\alpha_{n-t+1}, \beta_{n-t+1} \geq 1$. Let*

$$\begin{aligned} D^{\setminus} &= \{P_1, \dots, P_{j_r+1}, P_{j_r+2} + (1, -1), P_{j_r+3}, \dots, P_{n-t}, P_{n-t+1} + (-1, 1), P_{n-t+2}, \dots, P_n\}, \\ D^{\setminus\setminus} &= \{P_1, (0, 1), P_3, \dots, P_{n-t}, P_{n-t+1} + (1, -1), P_{n-t+2}, \dots, P_n\}. \end{aligned}$$

Then $2\Delta(D) \sim \Delta(D^{\setminus}) + \Delta(D^{\setminus\setminus})$ modulo $I_{<d}$ and staircase forms of lower partition types. Moreover, if the last block of $B(S)$ is not minimal or if $s_{n-t+1} > n - t_0$, then $\Delta(D) \sim \Delta(D^{\setminus\setminus})$ modulo $I_{<d}$ and staircase forms of lower partition types.

Proof. The reader is strongly recommended to see Example 30 first.

Suppose the partition type of D is

$$\underbrace{j_1 + \dots + j_1}_{m_1} + \dots + \underbrace{j_{r-1} + \dots + j_{r-1}}_{m_{r-1}} + \underbrace{j_r + \dots + j_r}_{m_r}.$$

Applying Lemma 26 to

$$\left(\sum x_i^{\alpha_{n-t+1}} y_i^{\beta_{n-t+1}-1} \right) \cdot \Delta(\{P_1, (0, 1), P_2, \dots, \widehat{P}_{n-t+1}, \dots, P_n\}),$$

which is an element in $I_{<d}$, we get a sum of n determinants: the 1st determinant is in $I_{<d}$ because the first row of its staircase form is the zero row. The 2nd determinant is

$$(3.1) \quad \Delta(\{P_1, P_{n-t+1}, P_2, \dots, \widehat{P}_{n-t+1}, \dots, P_n\}) = (-1)^{n-t-1} \Delta(D).$$

The i -th determinant for $i \geq 3$ is

$$\Delta(\{P_1, (0, 1), P_2, \dots, P_{i-2}, P_{i-1} + P_{n-t+1} - (0, 1), P_i, \dots, \widehat{P}_{n-t+1}, \dots, P_n\}),$$

when $3 \leq i \leq j_r + 3$, its partition type is equal to or lower than

$$\underbrace{j_1 + \dots + j_1}_{m_1} + \dots + \underbrace{j_{r-1} + \dots + j_{r-1}}_{m_{r-1}} + \underbrace{j_r + \dots + j_r}_{m_r-1} + (i-3) + (j_r - i + 3)$$

which is strictly lower than the partition type of D when $4 \leq i \leq j_r + 2$; when $i > j_r + 3$, the determinant is equivalent to 0. So modulo $I_{<d}$ and staircase forms of lower partition types, the sum of

$$(3.2) \quad \Delta(\{P_1, (0, 1), P_2 + P_{n-t+1} - (0, 1), P_3, \dots, \widehat{P}_{n-t+1}, \dots, P_n\}),$$

$$(3.3) \quad \Delta(\{P_1, (0, 1), P_2, \dots, P_{j_r+1}, P_{j_r+2} + P_{n-t+1} - (0, 1), P_{j_r+3}, \dots, \widehat{P}_{n-t+1}, \dots, P_n\})$$

and (3.1) is equivalent to 0.

Similarly as above, applying Lemma 26 to

$$\left(\sum x_i^{\alpha_{n-t+1}-1} y_i^{\beta_{n-t+1}}\right) \cdot \Delta(\{P_1, (0, 1), P_2, \dots, P_{j_r+1}, P_{j_r+2} + (1, -1), P_{j_r+3}, \dots, \widehat{P}_{n-t+1}, \dots, P_n\}),$$

which is an element in $I_{<d}$, we get a sum of n determinants: the 2nd determinant is

$$(3.4) \quad \Delta(\{P_1, P_{n-t+1} + (-1, 1), P_2, \dots, P_{j_r+1}, P_{j_r+2} + (1, -1), P_{j_r+3}, \dots, \widehat{P}_{n-t+1}, \dots, P_n\}),$$

the 3rd determinant is

$$(3.5) \quad \Delta(\{P_1, (0, 1), P_{n-t+1}, P_3, \dots, P_{j_r+1}, P_{j_r+2} + (1, -1), P_{j_r+3}, \dots, \widehat{P}_{n-t+1}, \dots, P_n\}),$$

the $(j_r + 3)$ -th determinant is

$$(3.6) \quad \Delta(\{P_1, (0, 1), P_2, \dots, P_{j_r+1}, P_{j_r+2} + P_{n-t+1} - (0, 1), P_{j_r+3}, \dots, \widehat{P}_{n-t+1}, \dots, P_n\}),$$

and all other determinants are equivalent to 0 modulo $I_{<d}$ and staircase forms of partition types lower than D . Now compare the two relations we obtained:

$$\begin{cases} (3.1) + (3.2) + (3.3) \sim 0, \\ (3.4) + (3.5) + (3.6) \sim 0. \end{cases}$$

Note that by Transfactor Lemma (Lemma 22), the polynomial (3.5) is equivalent to

$$\begin{aligned} & \Delta(\{P_1, P_2, P_{n-t+1}, P_3, \dots, P_{j_r+1}, P_{j_r+2}, P_{j_r+3}, \dots, \widehat{P}_{n-t+1}, \dots, P_n\}) \\ &= (-1)^{n-t-2} \Delta(D) = -(3.1), \end{aligned}$$

and also note that (3.3)=(3.6). So we have

$$(3.4) \sim -(3.5) - (3.6) \sim (3.1) - (3.3) \sim 2(3.1) + (3.2).$$

Since (3.4)= $(-1)^{n-t-1} \Delta(D^\setminus)$ and (3.2)= $(-1)^{n-t-2} \Delta(D^\setminus)$, the lemma follows.


Note that since $\deg P_{n-t+1} \geq \deg P_{n-t_0+1} = n - t_0$, we have

$$\deg(P_{j_r+2} + P_{n-t+1} - (0, 1)) \geq (j_r + 1) + (n - t_0) - 1 = j_r + n - t_0$$


which is greater than $n - 1$ if $j_r \geq t_0$. But this is always the case if the last block of $B(S)$ is not minimal. In this case, (3.3) ~ 0 and therefore (3.1) + (3.2) ~ 0 . Of course we still have (3.1) + (3.2) ~ 0 if $(s_{n-t+1}) \deg P_{n-t+1} > n - t_0$. $\square$

The discovery of Lemma 29 is motivated by the observation in the following example.

Example 30. Let $n = 9$, $k = 3$, then $d = \binom{9}{2} - 3 = 33$. Consider a partition $(2 + 1) \in \Pi_3$,



and let $t = 3$. Let $D =$ (in the standard order) and $f = \Delta(D)$.



(i) Applying Lemma 26 to the product of $\sum_{i=1}^9 x_i^5 y_i^0$ with $\Delta(\dots)$. Modulo $I_{<d} +$ (minimal staircase forms of lower partitions), there are 2 summands remained in the

sum: $\Delta(\text{grid1})$ and $\Delta(\text{grid2})$. Since the former is $\pm f$, the latter is equivalent to $\pm f$ modulo $I_{<d} + (\text{minimal staircase forms of lower partitions})$.

(ii) On the other hand, by Transfactor Lemma, $\Delta(\text{grid3})$ is in $I_{<d} + (\text{minimal staircase forms of lower partitions}) + (f)$. Note that we move the points $(1, 0)$ and $(0, 3)$ in D .

(iii) Applying Lemma 26 to the product of $\sum_{i=1}^9 x_i^4 y_i^1$ with $\Delta(\text{grid4})$. Modulo $I_{<d} + (\text{minimal staircase forms of lower partitions})$, we have 3 summands left: $\Delta(\text{grid5})$, $\Delta(\text{grid6})$, and $\Delta(\text{grid7})$. We already know that the first two are in the ideal $I_{<d} + (\text{minimal staircase forms of lower partitions}) + (f)$, hence the last one as well. $\square$

Proof of Proposition 20. First we explain the condition $n \geq 8k + 5$. It follows from the conditions $d_1 + d_2 \leq n(n-1)/2$ and $d_1, d_2 \geq (2k+1)n$, which imply $n(n-1)/2 \geq 2(2k+1)n$, equivalently $n \geq 8k + 5$.

We prove by induction on k . The base case $k = 0$ is proved in Lemma 23. Suppose the proposition is proved for $< k$.

Let $D = \{P_1, \dots, P_n\} \in \mathfrak{D}$, S be a minimal staircase form of D of partition type μ . Notice that, without loss of generality, we can assume that the last block of $B(S)$ is of size greater than 1. Indeed, suppose the last block, which corresponds to P_n , is of size 1, and suppose that the block M is the last block among those of size greater than 1. Since $d_1 \geq (2k+1)n$, there are sufficient size-1 blocks in $B(S)$, such that by successively moving a P_i corresponding to a size-1 block to northwest direction and moving P_n to southeast direction using Transfactor Lemma, we can assume $P_n = (\alpha_n, 0)$. (Of course $d_1 \geq (2k+1)n$ is not a sharp bound. We obtain this bound by noticing that there are at most $2k$ points of D that do not correspond to size-1 blocks, the x -degree of each of which is less than n , while the last point P_n also has x -degree less than n . So as long as the total x -degree is larger than $2k \cdot n + n$, the point P_n can be moved to southeast direction by Transfactor lemma.) Then we can apply Minors Permuting Lemma to permute the last block with the blocks before it until it moves in front of M . Then M is moved to the lower right in a block diagonal form. This procedure can be repeated until M becomes the last block.

By Transfactor Lemma and Minors Permuting Lemma together with the condition that $n \geq 8k + 5$, we can assume the first $(k + 2)$ blocks of $B(S)$ are all of size 1.

Now we are in the position to apply Lemma 29. Denote by t_0 the size of the last block in $B(S)$. By Transfactor Lemma we may assume $P_2 = (1, 0)$. If for $1 \leq t \leq t_0$ the point P_{n-t+1} has degree $s_{n-t+1} > n - t_0$, then $D \sim D^{\searrow}$, which means that we can move P_{n-t+1} to $P_{n-t+1} + (1, -1)$. Successively applying this procedure, we may assume that all points P_i for $i > n - t + 2$ have y-coordinates 0.

Define $a(D) = \alpha_{n-t_0+2} - \alpha_{n-t_0+1}$. Then

$$a(D^{\nwarrow}) - 1 = a(D) = a(D^{\searrow}) + 1.$$

Consider the special case when $P_{n-t_0+1} = P_{n-t_0+2}$. In this case $\Delta(D) = 0$ hence $\Delta(D^{\nwarrow}) \sim -\Delta(D^{\searrow})$, $a(D^{\nwarrow}) = 1$ and $a(D^{\searrow}) = -1$. Let D'' be the set obtained by interchanging the $(n - t_0 + 1)$ -th and $(n - t_0 + 2)$ -th points in $D^{\searrow}$. Now we compare $D^{\nwarrow} = \{P'_1, \dots, P'_n\}$ with $D'' = \{P''_1, \dots, P''_n\}$:

- They both give minimal staircase forms with the same partition type as S ,
- $a(D^{\nwarrow}) = a(D'') = 1$,
- $\Delta(D^{\nwarrow}) \sim \Delta(D'')$,
- $P''_i = \begin{cases} P'_i + (1, -1), & \text{for } i = n - t + 1, n - t + 2; \\ P'_i + (-1, 1), & \text{for } i = 2, j_r + 2; \\ P'_i, & \text{otherwise.} \end{cases}$

In other words, we can move P'_{n-t+1} and P'_{n-t+2} of $D^{\nwarrow}$ to southeast direction and move two size-1 blocks of $D^{\nwarrow}$ to northwest direction simultaneously without changing $\Delta(D^{\nwarrow})$ modulo the equivalence relation. Repeat the procedure until the y-coordinates of the $(n - t + 1)$ -th and $(n - t + 2)$ -th points are 1 and 0, respectively. Then apply the inductive assumption for the first $n - t$ points, we can draw the following conclusion:

For any D' and D'' such that

- (i) both have minimal staircase forms,
- (ii) their staircase forms are of the same partition type,
- (iii) $\Delta(D)$ and $\Delta(D')$ have the same bi-degree,
- (iv) $a(D') = a(D'') = 1$,

then $\Delta(D') \sim \pm \Delta(D'')$. If (ii) is replaced by a stronger condition:

- (ii)' they are both in standard order and their block diagonal forms are of the same shape (i.e. for any i , the size of the i -th blocks in both block diagonal forms are the same),

then $\Delta(D') \sim \Delta(D'')$.

By Lemma 29, we can also show that, under condition (i) (ii)' (iii) and assume that $a(D'), a(D'') > 0$,

$$\Delta(D')/a(D') \sim \Delta(D'')/a(D'').$$

Indeed, it is sufficient to show that

$$(3.7) \quad \text{if conditions (i)(ii)' (iii) hold and } a(D') = 1, \text{ then } a(D'')\Delta(D') \sim \Delta(D'').$$

This can be proved by induction on $a(D'')$. The case $a(D'') = 0$ is trivial since in this case $\Delta(D'') = 0$. We have already shown the case $a(D'') = 1$. Now by inductive assumption we assume that (3.7) is true for $a(D'') = k - 1$ and k . Suppose $a(D'') = k + 1$. Take $D \in \mathfrak{D}$ such that $D^{\nwarrow} \sim D''$. (This is always possible by using Transfactor Lemma and Minors Permuting Lemma to modify D'' .) Then Lemma 29 asserts that $2\Delta(D) \sim \Delta(D^{\nwarrow}) + \Delta(D^{\searrow})$. By inductive assumption $\Delta(D) \sim k\Delta(D')$ and $\Delta(D^{\searrow}) \sim (k - 1)\Delta(D')$, therefore

$$\Delta(D'') \sim \Delta(D^{\nwarrow}) \sim 2k\Delta(D') - (k - 1)\Delta(D') = (k + 1)\Delta(D'),$$

this completes the inductive proof of (3.7).

As an immediate consequence, any minimal staircase form of partition type μ generates all the minimal staircase forms of the same partition type μ , modulo $I_{<d} + (\text{minimal staircase forms of partition type } <_P \mu)$. This completes the proof. $\square$

Now we can prove Proposition 19.

Proof of Proposition 19. Assume $D = \{P_1, \dots, P_n\} \in \mathfrak{D}$ and S is a staircase form of D and is not minimal. By Transfactor Lemma and Minors Permuting Lemma, we can assume without loss of generality that, in the block diagonal form $B(S) = \text{diag}(B_1, \dots, B_s)$, all the size-1 blocks stand before the blocks of size greater than 1.

First note that if the assumption of Lemma 29 is satisfied and the last block of $B(S)$ is not minimal, the conclusion easily follows. Indeed, in this case the claim $\Delta(D) \sim \Delta(D')$ in Lemma 29 implies that we may move any point P_i in the last block of $B(S)$ to $P_i + (1, -1)$. Start from a point P_i for some i that $n - t_0 + 1 \leq i \leq n - 1$ such that it has the same degree as P_{i+1} . Keep on moving P_i to southeast direction until it collides with P_{i+1} and then the determinant will be 0.

Now we show that we can always assume the assumption of Lemma 29 is satisfied and the last block of $B(S)$ is not minimal. The assumption of Lemma 29 is always satisfied by using Minors Permuting Lemma and Transfactor Lemma, since there are sufficient size-1 blocks in $B(S)$. To finish the proof of the proposition, we only need to exclude the case when the last block B_s of $B(S)$ is minimal. Denote the size of B_s by $t_0 \geq 2$. Define D_1 to be the set $\{P_1, \dots, P_{n-t_0}\}$, define $n' = n - t_0$, and let d', d'_1, d'_2, k' be the total degree, x-degree, y-degree and deficit of D_1 , respectively. Then $k \geq k' + t_0 - 1$ so

$$\begin{aligned} n' &\geq 8k + 5 - t_0 \geq 8(k' + t_0 - 1) + 5 - t_0 \geq 8k' + 5, \\ d'_1 &> d_1 - t_0 n \geq (2k + 1)n - t_0 n \geq (2k' + t_0 - 1)n \geq (2k' + 1)n' \end{aligned}$$

and similarly $d'_2 \geq (2k' + 1)n'$. Then we can use induction to assert that $\Delta(D_1)$ is generated by elements in $I_{<d'}$ and minimal staircase form of degree d' . Now $\Delta(D) = \Delta(D_1) \cdot \det(B_s)$ is generated by $I_{<d}$ and minimal staircase form of degree d . Hence in the case when B_s is minimal, there is nothing to prove. $\square$

In order to complete the proof of Theorem 6, we use the following Lemma. Recall that

$$a(\lambda) = \sum_i (n - i - \lambda_i),$$

$$b(\lambda) = \#\{i < j : \lambda_i - \lambda_j + i - j \in \{0, 1\}\}.$$

Lemma 31. *Let u be an positive integer that $k \leq u \leq n - 2$ and let $v = n - 1 - u$. If $d_1 = u(u + 1)/2$ and $d_2 = v(v + 1)/2 + uv - k$, then $\dim M_{d_1, d_2} \geq p(k)$.*

Proof. Consider a partition

$$\lambda = (u + \varepsilon_0, u - 1 + \varepsilon_1, u - 2 + \varepsilon_2, \dots, 1 + \varepsilon_{u-1}, 0, 0, \dots, 0),$$

where $(v + 1)$ zeroes are at the end. If

$$(3.8) \quad \begin{cases} \varepsilon_i = 0 \text{ or } 1 \text{ for } 0 \leq i \leq u - 1, \\ \sum_{i=0}^{u-1} \varepsilon_i = k, \\ \text{and } \sum_{i=0}^{u-1} i\varepsilon_i = k(k + 1)/2, \end{cases}$$

then it is straightforward to check that λ satisfies

$$b(\lambda) = u(u + 1)/2, \text{ and } a(\lambda) = v(v + 1)/2 + uv - k.$$

Since there are $p(k)$ number of solutions for the system (3.8), we have $\dim M_{d_1, d_2} \geq p(k)$. $\square$

Now we are ready to prove Theorem 6.

Proof of Theorem 6. Corollary 21 asserts that M_{d_1, d_2} is generated by $p(k)$ elements

$$\{\det(S_\mu)\}_{\mu \in \Pi_k}$$

where S_μ is an arbitrary minimal staircase form of bidegree (d_1, d_2) and of partition type μ . So to complete the proof, we need to show the minimality of the above generators, which is equivalent to show $\dim M_{d_1, d_2} \geq p(k)$. Lemma 31 provides such a lower bound of $\dim M_{d_1, d_2}$ for special values of d_1 and d_2 . For general values of d_1 and d_2 , the idea is to add sufficiently many appropriate size-1 blocks such that we can apply Lemma 31. We shall explain as below.

Choose a sufficiently large number $\tilde{n} \gg n$ such that there are positive integers u and v satisfying $k \leq u \leq \tilde{n} - 2$, $1 + u + v = \tilde{n}$, $u(u + 1)/2 \geq (2k + 1)\tilde{n}$, and $v(v + 1)/2 + uv - k \geq (2k + 1)\tilde{n}$. Choose $(\tilde{n} - n)$ points $P_i = (\alpha_i, \beta_i)$ for $n + 1 \leq i \leq \tilde{n}$ so that

$$\alpha_i + \beta_i = i - 1 \quad (n + 1 \leq i \leq \tilde{n}),$$

$$\sum_{i=1}^{\tilde{n}} \alpha_i = u(u + 1)/2 =: \tilde{d}_1,$$

$$\sum_{i=1}^{\tilde{n}} \beta_i = v(v + 1)/2 + uv - k =: \tilde{d}_2$$

which is always possible. By our choice of P_i ($n+1 \leq i \leq \tilde{n}$), if $D = \{P_1, \dots, P_n\}$ has a minimal staircase form of partition type μ , then $\tilde{D} = \{P_1, \dots, P_n, P_{n+1}, \dots, P_{\tilde{n}}\}$ also has a minimal staircase form of the same partition type μ . Let $\tilde{S}$ be the staircase form of $\tilde{D}$ and $B(\tilde{S})$ the block diagonal form of $\tilde{S}$. Denote by f_0 the product of the last $(\tilde{n} - n)$ size-1 minors in $B(\tilde{S})$. Let $\tilde{I} = \cap_{1 \leq i < j \leq \tilde{n}} (x_i - x_j, y_i - y_j)$ be an ideal of $\mathbb{C}[x_1, y_1, \dots, x_{\tilde{n}}, y_{\tilde{n}}]$, define $\tilde{M} = \tilde{I}/(\mathbf{x}, \mathbf{y})\tilde{I}$ which is doubly graded as $\oplus_{\tilde{d}_1, \tilde{d}_2} \tilde{M}_{\tilde{d}_1, \tilde{d}_2}$. Then we have a $\mathbb{C}$ -linear map:

$$\begin{aligned} L : M_{d_1, d_2} &\rightarrow \tilde{M}_{\tilde{d}_1, \tilde{d}_2} \\ f &\mapsto f \cdot f_0. \end{aligned}$$

For every partition μ of k , $L(\det S_\mu)$ is of partition type μ . Since $\{L(\det S_\mu)\}_{\mu \in \Pi(k)}$ form a basis for $\tilde{M}_{\tilde{d}_1, \tilde{d}_2}$, the map L is surjective. Therefore $\dim M_{d_1, d_2} \geq \dim \tilde{M}_{\tilde{d}_1, \tilde{d}_2} \geq p(k)$, which provides the expected lower bound for $\dim M_{d_1, d_2}$. $\square$

4. CONJECTURAL SET OF GENERATORS

Recall that Λ is the set of integer sequences $\lambda_1 \geq \dots \lambda_{n-1} \geq \lambda_n = 0$ satisfying $\lambda_i \leq n - i$ for all i . We propose the following conjecture.

Conjecture 32. *For any $\lambda \in \Lambda$, let*

$$a_i = n - i - \lambda_i, \quad b_i = \#\{i < j : \lambda_i - \lambda_j + i - j \in \{0, 1\}\}$$

and $D(\lambda) = \{(a_i, b_i) | 1 \leq i \leq n\}$. Then $G := \{\Delta(D(\lambda))\}_{\lambda \in \Lambda}$ generates I .

The cases for $n \leq 8$ have been verified by computer. In our forthcoming paper, we will show that this conjecture holds true for certain bi-degree spaces M_{d_1, d_2} .

Remark 33. An equivalent conjecture is given by Mahir Can and Nick Loehr in their unpublished work.

REFERENCES

- [1] N. Bergeron and Z. Chen, Basis of Diagonally Alternating Harmonic Polynomials for low degree, arXiv: 0905.0377.
- [2] A. M. Garsia and J. Haglund, A positivity result in the theory of Macdonald polynomials, Proc. Natl. Acad. Sci. USA **98** (2001), no. 8, 4313–4316 (electronic).
- [3] A. M. Garsia and J. Haglund, A proof of the q, t-Catalan positivity conjecture, Discrete Math. **256** (2002), no. 3, 677717, LaCIM 2000 Conference on Combinatorics, Computer Science and Applications (Montreal, QC).
- [4] Mark Haiman, Hilbert schemes, polygraphs and the Macdonald positivity conjecture, J. Amer. Math. Soc. **14** (2001), no. 4, 941–1006.
- [5] Mark Haiman, Vanishing theorems and character formulas for the Hilbert scheme of points in the plane, Invent. Math. **149** (2002), no. 2, 371–407.
- [6] Mark Haiman, Commutative algebra of n points in the plane, With an appendix by Ezra Miller. Math. Sci. Res. Inst. Publ., 51, Trends in commutative algebra, 153–180, Cambridge Univ. Press, Cambridge, 2004.
- [7] G.H. Hardy, Ramanujan: twelve lectures on subjects suggested by his life and work, Chelsea Publishing Company, New York 1959.

- [8] K. Lee and L. Li, q, t -Catalan numbers and generators for the radical ideal defining the diagonal locus of $(\mathbb{C}^2)^n$.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907

E-mail address: `kyungl@purdue.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA CHAMPAIGN, URBANA, IL 61801

E-mail address: `llpku@math.uiuc.edu`